Check List the journal of biodiversity data
NOTES ON GEOGRAPHIC DISTRIBUTION

First record of *Bichromomyia flaviscutellata* (Mangabeira, 1942) (Diptera, Psychodidae, Phlebotominae), a vector of *Leishmania* (*Leishmania*) *amazonensis* (Lainson & Shaw, 1972), in the state of Paraíba, Brazil

Lucas José Macêdo Freire[1], Alessandre Pereira-Colavite[2], Maurício Luiz Vilela[1]

1 *Laboratório Interdisciplinar de Vigilância em Diptera e Hemiptera, Instituto Oswaldo Cruz, Fiocruz, Rio de Janeiro, Rio de Janeiro, Brazil*
2 *Laboratório de Entomologia, Departamento de Sistemática e Ecologia, CCEN, Universidade Federal da Paraíba, João Pessoa, Paraíba, Brazil*

Corresponding author: *Lucas José Macêdo Freire (macedolucasjf@gmail.com)*

Abstract. *Bichromomyia flaviscutellata* (Mangabeira 1942), a vector of *Leishmania* (*Leishmania*) *amazonensis* (Lainson & Shaw 1972), has been recorded for the first time in the state of Paraíba, northeastern Brazil. A female specimen was captured using an HP-type light trap in an urban park in the municipality of João Pessoa in December 2022. The new record of Bi. flaviscutellata enhances our understanding of the sandfly fauna in Paraíba, particularly in an epidemiological context, given its significance in the transmission of Leishmania pathogens.

Key words. Atlantic Forest, geographic distribution, leishmaniasis, sandfly

Freire LJM, Pereira-Colavite A, Vilela ML (2025) First record of *Bichromomyia flaviscutellata* (Mangabeira, 1942) (Diptera, Psychodidae, Phlebotominae), a vector of *Leishmania* (*Leishmania*) *amazonensis* (Lainson & Shaw, 1972), in the state of Paraíba, Brazil. Check List 21 (2): 368–373. https://doi.org/10.15560/21.2.368

INTRODUCTION

Sandflies, blood-feeding insects belonging to the subfamily Phlebotominae (Diptera, Psychodidae), are recognized vectors of pathogens that cause diseases, including viruses, bacteria, and most notably, protozoa of the genus *Leishmania* Ross, 1903 (Brazil and Brazil 2003). Currently, approximately 1,060 species of sandflies have been identified worldwide, with 539 species found in the Americas. Hosting 279 species, Brazil is among the countries with the highest diversity of sandflies (de Aguiar and Vieira 2018; Galati 2023). About 10% of these species are vectors of the infectious agents responsible for leishmaniasis (Rangel and Lainson 2009; Shaw et al. 2018).

Leishmaniasis in Brazil, initially associated with rural and forested environments, has undergone urbanization due to human intervention in the environment and the adaptability of sandflies. This has led to a higher incidence of the disease in domestic and peri-domestic areas within urban centers (Harhay et al. 2011; Temponi et al. 2018). Additionally, the proximity of human dwellings to forest fragments and the rise in ecotourism have increased the risk of human contact with vectors, thereby facilitating the transmission of leishmaniasis (Uzcategui et al. 2020; Thomaz-Soccol et al. 2023). Understanding the distribution and composition of these insect species is therefore crucial to understanding the dynamics of leishmaniasis in areas both with and without reported cases of the disease.

In Brazil, there are two primary forms of leishmaniasis: American visceral leishmaniasis (AVL), which is considered the most fatal, and American tegumentary leishmaniasis (ATL), which can manifest in cutaneous, mucocutaneous, or diffuse forms (Maroli et al. 2013). In the Brazilian Northeast, although the incidence of AVL is high, the number of cases has remained stable (Bruhn et al. 2023). The region also reports many cases of ATL, despite having one of the lowest diversities of sandflies compared to other regions (de Aguiar and Vieira 2018). In Paraíba, there is limited recorded diversity of these insects, and few studies have been conducted on their fauna, leading to undersampling (Silva et al. 2022), even in municipalities with endemic cases of leishmaniasis (SINAN 2023). This study aims to document for the first time the presence of the species *Bichromomyia flaviscutellata* (Mangabeira 1942), a vector of *Leishmania* (*Leishmania*) *amazonensis*

OPEN ACCESS

Academic editor: Fabio Laurindo da Silva
Received: 13 September 2024
Accepted: 11 November 2024
Published: 1 April 2025

Copyright © The authors. This is an open-access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0)

(Lainson & Shaw 1972), in Paraíba, thereby contributing to the knowledge of sandfly fauna in the state. The motivation or purpose of your research should appear in the Introduction, providing some of the historical basis for those questions.

METHODS

Monthly collections were conducted at the Monitoring Station of Cuiá Municipal Natural Park (EMPNMC) (07°11′27.0″S, 034°51′23.4″W) under permanent license SISBIO No. 81583-1. Five CDC light traps (HP model) (Pugedo et al. 2005) were placed along the park's main trails in areas likely to attract sandflies, such as tree hollows, sites with high organic matter content, and animal burrows. The traps were active for three consecutive nights each month, from 5:00 PM to 7:00 AM, from May 2022 to May 2024.

The captured insects were transported to the laboratory in nylon nets, where they were euthanized by cryogenic methods, sorted, and stored in tubes containing 70% alcohol. The specimens then underwent clarification and diaphanization processes, which included immersion in 10% KOH for 3 h, followed by absolute acetic acid for 20 min, a 30-min rinse in distilled water (Type II), and a 24-h immersion in lactophenol to facilitate the identification of taxonomic structures (Vilela et al. 2018).

Specimens were mounted on slides using Berlese solution and examined under a stereomicroscope. Identification was conducted using an optical microscope with the aid of Galati (2023) taxonomic key and the species abbreviations proposed by Marcondes (2007). All specimens were deposited in the Entomological Collection of the Federal University of Paraíba (DSEC). Thematic maps were created using QGIS Desktop v. 3.36.0 (QGIS 2024), and specimen photographs were taken with a Carl Zeiss Primostar binocular microscope equipped with an integrated camera.

RESULTS

During the 25 months of study, a single female *Bichromomyia flaviscutellata* (Mangabeira 1942) was captured in December 2022 at the Monitoring Station of Cuiá Municipal Natural Park (EMPNMC) (Figure 1), specifically at the capture site (Figure 1, red dot) located in a hollow tree trunk in a forested area.



Figure 1. Capture point where the species *Bichromomyia flaviscutellata* was recorded at the Monitoring Station of the Cuiá Municipal Natural Park (EMPNMC) during the months of May 2022 to May 2024 in the municipality of João Pessoa, Paraíba.



Figure 2. Specimen of *Bichromomyia flaviscutellata* collected in João Pessoa, Paraíba. **A.** Ventral view of the head; arrow showing the clypeus. **B.** Ventral view of the abdomen with the spermatheca of the female; arrow showing the relationship between the common spermathecal duct and the stem region of the genital furca. **C.** Lateral view of the thorax; arrow showing the bicolor division of the thorax. **D.** Ventral view of the head; arrow showing the anterior teeth of the cibarium. Scale bar: A = 100 µm; B = 10 µm; C = 100 µm; D = 10 µm.

***Bichromomyia flaviscutellata* (Mangabeira, 1942)**
Figure 2

New record. BRAZIL – PARAÍBA • João Pessoa, Cuiá Municipal Natural Park, Capture Point 3; 07°11′27.3″S, 034°51′27.9″W; 112 m alt; 22.XII.2022; L. J. Macedo, collector; HP light trap, hollow tree trunk; 1 ♀, slide mounted, DSEC005499DP.

Identification. A single individual of the species was identified. The key characteristics include a clypeus equal in size to the eyes (Figure 2A) and a bicolored mesonotum, with the posterior part of the scutum and scutellum lighter than the rest, which is dark brown (Figure 2C). Interspecifically, the spermatheca of the female *Bi. flaviscutellata* differs from that of *Bi. reducta* (Feliciangeli, Ramirez-Pérez & Ramirez, 1988) by the width of the common duct relative to the median region of the genital furca, which is narrower in the former species (Figure 2B). The apical ring is narrower than the pre-apical ring, with a head larger than the sum of the last three rings, oriented toward the center of the genital furca (Figure 3B). The cibarial region of *Bi. flaviscutellata* contains 6–8 horizontal teeth (Figure 2D).

DISCUSSION

The genus *Bichromomyia* (Galati, 2003) includes four species, among them, three subspecies of *Bi. olmeca*, distributed throughout Neotropical region, primarily in the Amazon region. In Brazil, the species recorded are *Bichromomyia flaviscutellata*, *Bi. inornata* (Martins, Falcão & Silva, 1965), *Bi. olmeca* (Vargas & Díaz-Nájera, 1959) (ssp. *bicolor* (Fairchild & Theodor, 1971) and *Bi. nociva* (Young & Arias, 1982)), and *Bi. reducta*. Of these, *Bi. flaviscutellata*, *Bi. olmeca nociva*, and *Bi. reducta* are epidemiologically significant as potential vectors of *L*. (*L*.) *amazonensis*, the causative agent of ATL in its diffuse form (Rangel et al. 2018).

Bichromomyia flaviscutellata was first described by Mangabeira in 1942, based on a male specimen collected in Belém, Pará. This species has an almost cosmopolitan distribution in Brazil, present in 19 of the 26 states and the Federal District, in all of the five Brazilian regions (Costa et al. 2009), and it is also found in other countries within the Amazon region (Figure 3). Records of *Bi. flaviscutellata* are common in the Northern, Northeastern, Central-Western, and Southeastern regions of Brazil, across diverse habitats ranging from primary forests to peri-urban areas (Andrade et al. 2009; Vilela et al. 2011; Carvalho et al. 2013; de Aguiar and Vieira, 2018; Rangel et al. 2018; Galati 2023).

Documenting *Bi. flaviscutellata* in a new state broadens our understanding of the eco-epidemiology of leishmaniasis, enhancing our knowledge of the distribution and ecology of this vector species. Despite its highly rodentophilic behavior (Lainson and Shaw 1968) and low attraction to humans, *Bi. flaviscutellata* is the primary vector of *L*. (*L*.) *amazonensis* in Brazil (Lainson and Shaw 1972), which causes ATL in its diffuse form. Environmental and climatic changes in Brazil may be contributing to the expansion of this species into peri-urban areas, thereby increasing the incidence of new leishmaniasis cases (Shaw et al. 2007; Queiroz et



Figure 3. Geographic distribution map of *Bichromomyia flaviscutellata* in South America. Brazil highlighted (in dark green) with its political division into states. Abbreviations of the Brazilian states (acronyms not underlined): AC, Acre; AL, Alagoas; AP, Amapá; AM, Amazonas; BA, Bahia; CE, Ceará; DF, Federal District; ES, Espírito Santo; GO, Goiás; MA, Maranhão; MT, Mato Grosso; MS, Mato Grosso do Sul; MG, Minas Gerais; PA, Pará; PB, Paraíba; PR, Paraná; PE, Pernambuco; PI, Piauí; RJ, Rio de Janeiro; RN, Rio Grande do Norte; RS, Rio Grande do Sul; RO, Rondônia; RR, Roraima; SC, Santa Catarina; SP, São Paulo; SE, Sergipe; TO, Tocantins. Abbreviations of the South American countries (light green) (underlined acronyms): ARG, Argentina; BOL, Bolivia; CHL, Chile; COL, Colombia; ECU, Ecuador; GUF, French Guiana; GUY, Guyana; PRY, Paraguay; PER, Peru; SUR, Suriname; URY, Uruguay; VEN, Venezuela; TRT, Trinidad and Tobago. Brazilian limits in bold.

al. 2012). Carvalho et al. (2015) observed that in a scenario with environmental changes, through ecological niche modeling algorithms, the range of environments suitable areas for *Bi. flaviscutellata* will be expanded throughout Brazil, especially in areas where human density would be greater than today.

The capture of *Bi. flaviscutellata* in this study occurred in a fragment of the Atlantic Forest used for ecotourism. This suggests a possible interaction between humans and the enzootic cycle of leishmaniasis, potentially leading to infection (Brilhante et al. 2015; Fonteles et al. 2018). Additionally, this forest fragment is in a recently urbanized area experiencing significant human pressure, which could facilitate the spread of this species into peri-urban areas (Queiroz et al. 2012). This finding opens up new opportunities for research on the bioecology of this vector and molecular analyses for the identification of Leishmania, aiming to improve leishmaniasis knowledge and surveillance.

With the new record of *Bi. flaviscutellata*, the number of sandfly species in Paraíba increases from 16 to 17, although the state remains one of those with the lowest species richness in the Northeastern Region (de Aguiar and Vieira 2018; Silva et al. 2022).

ACKNOWLEDGEMENTS

We are grateful to the Laboratório de Entomologia da Universidade Federal da Paraíba (LabEnt) and the Laboratório Interdisciplinar de Vigilância Entomológica em Diptera e Hemiptera (LIVEDIH) for providing the appropriate infrastructure for the capture, processing and identification processes of the specimens and to the members for the company during the fields survey. We would like to thank Francisco do Cuiá for helping during the surveys. We extend our profound gratitude to the academic editor and reviewers of this work for their invaluable contributions, which have significantly enhanced the quality of the manuscript.

ADDITIONAL INFORMATION

Conflict of interest

The authors declare that no competing interests exist.

Ethical statement

No ethical statement is reported.

Funding

This study was financially supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001.

Author contributions

Conceptualization: LJMF, MLV. Data curation: LJMF. Formal analysis: LJMF. Investigation: LJMF. Methodology: LJMF, MLV. Resources: LJMF, APC. Supervision: APC, MLV. Validation: MLV. Visualization: LJMF, APC. Writing – original draft: LJMF. Writing – review and editing: APC, MLV.

Author ORCID iDs

Lucas José Macêdo Freire https://orcid.org/0000-0003-3243-2847
Alessandre Pereira-Colavite https://orcid.org/0000-0002-7660-8384
Maurício Luiz Vilela https://orcid.org/0000-0001-7610-0719

Data availability

All data that support the findings of this study are available in the main text.

REFERENCES

Andrade ARO, Nunes VLB, Galati EAB, Arruda CCP, Santos MFC, Rocca MEG, Aquino RB (2009) Epidemiological study on leishmaniasis in an area of environmental tourism and ecotourism, State of Mato Grosso do Sul, 2006–2007. Revista da Sociedade Brasileira de Medicina Tropical 42 (5): 488–493: https://doi.org/10.1590/S0037-86822009000500003

Brazil RP, Brazil BG (2003) Biologia de flebotomíneos Neotropicais. In: Rangel EF, Lainson R (Eds.) Flebotomíneos do Brasil. Editora da Fundação Oswaldo Cruz, Rio de Janeiro, Brazil, 257–274.

Brilhante AF, Nunes VL, Kohatsu KA, Galati EA, Rocca ME, Ishikawa EA (2015) Natural infection of phlebotomines (Diptera: Psychodidae) by *Leishmania* (*Leishmania*) *amazonensis* in de aguian area of ecotourism in Central-Western Brazil. Journal of Venomous Animals and Toxins Including Tropical Diseases 21 (39): 1–3: https://doi.org/10.1186/s40409-015-0041-8

Bruhn FRP, Werneck GL, Barbosa DS, Câmara DCP, Simões TC, Buzanovsky LP, Duarte AGS, de Melo SN, Cardoso DT, Donato LE, Maia-Elkhoury ANS, Belo VS (2023) Spatio-temporal dynamics of visceral leishmaniasis in Brazil: A nonlinear regression analysis. Zoonoses and Public Health 71: 144–156: https://doi.org/10.1111/zph.13092

Carvalho BM, Rangel EF, Ready PD, Vale MM (2015) Ecological niche modelling predicts southward expansion of *Lutzomyia* (*Nyssomyia*) *flaviscutellata* (Diptera: Psychodidae: Phlebotominae), vector of *Leishmania* (*Leishmania*) *amazonensis* in South America, under Climate Change. PLoS ONE 10 (11): e0143282. https://doi.org/10.1371/journal.pone.0143282

Carvalho BM, Maximo M, Costa WA, Santana ALF, Costa, SM, Rego TANC, Pita-Pereira D, Rangel EF (2013) Leishmaniasis transmission in an ecotourism area: potential vectors in Ilha Grande, Rio de Janeiro State, Brazil. Parasites & Vectors 6 (325): 1–12. https://doi.org/10.1186/1756-3305-6-325

Costa JML, Costa AAUML, Elkhoury AN, Bezerril ACR, Barral A, Saldanha ACR (2009) Leishmaniose cutânea difusa (LCD) no Brasil após 60 anos de sua primeira descrição. Gazeta Médica da Bahia 79: 16–24.

de Aguiar GM, Vieira VR (2018) Regional distribution and habitats of Brazilian Phlebotomine species. In: Rangel EF, Shaw JJ (Eds.) Brazilian sand flies. Springer, Cham, Switzerland & Berlin, Germany, 251–298. https://doi.org/10.1007/978-3-319-75544-1_4

Fonteles RS, Pereira-Filho AA, Moraes JLP, Pereira SRF, Rodrigues BL, Rêbelo JMM (2018) Detection of *Leishmania* DNA and blood meal identification in sand flies (Diptera: Psychodidae) from Lençois Maranhenses National Park region, Brazil. Journal of Medical Entomology 55 (2): 445–451. https://doi.org/10.1093/jme/tjx230

Galati EAB (2023) Morfologia e terminologia de Phlebotominae (Diptera: Psychodidae). Classificação e identificação de táxons das Américas. Vol I. Apostila da disciplina bioecologia e identificação de Phlebotominae do programa de pós-graduação em saúde pública. Faculdade de Saúde Pública da Universidade de São Paulo, São Paulo, Brazil. https://www.fsp.usp.br/egalati/index.php/2018/07/24/materiais/. Accessed on: 2023-09-01.

Harhay MO, Olliaro PL, Costa DL, Costa CH (2011) Urban parasitology: visceral leishmaniasis in Brazil. Trends in Parasitology 27 (9): 403–409: https://doi.org/10.1016/j.pt.2011.04.001

Lainson R, Shaw JJ (1968) Leishmaniasis in Brazil: I. Observations on enzootic rodent leishmaniasis—incrimination of *Lutzomyia flaviscutellata* (Mangabeira) as the vector in the Lower Amazonian Basin. Transactions of the Royal Society of Tropical Medicine and Hygiene 62 (3): 385–395. https://doi.org/10.1016/0035-9203(68)90090-4

Lainson R, Shaw JJ (1972) Leishmaniasis in Brazil: VI. Observations on the seasonal variations of *Lutzomyia flaviscutellata* in different types of forest and its relationship to enzootic rodent leishmaniasis (*Leishmania mexicana amazonensis*). Transactions of the Royal Society of Tropical Medicine and Hygiene 66 (5): 709–171. https://doi.org/10.1016/0035-9203(72)90084-3

Marcondes CB (2007) A proposal of generic and subgeneric abbreviations for Phlebotomine Sandflies (Diptera: Psychodidae: Phlebotominae) of the world. Entomological News 118 (4): 351–356. https://doi.org/10.3157/0013-872x(2007)118[351:apogas]2.0.co;2

Maroli M, Feliciangeli MD, Bichaud L, Charrel RN, Gradoni L (2013) Phlebotomine sandflies and the spreading of leishmaniases and other diseases of public health concern. Medical and Veterinary Entomology 27: 123–147. https://doi.org/10.1111/j.1365-2915.2012.01034.x

Pugedo HR, Barata A, França-Silva AJ, Silva JC, Dias ES (2005) HP: an improved model of suction light trap for the capture of small insects. Revista da Sociedade Brasileira de Medicina Tropical 38: 70–72. https://doi.org/10.1590/s0037-86822005000100017

QGIS Development Team (2024) QGIS Geographic Information System. Open Source Geospatial Foundation Project. http://qgis.osgeo.org. Accessed on: 2024-8-15.

Queiroz MFM, Varjão JR, Moraes SCD, Salcedo GE (2012) Analysis of sandflies (Diptera: Psychodidae) in Barra do Garças, state of Mato Grosso, Brazil, and the influence of environmental variables on the vector density of *Lutzomyia longipalpis* (Lutz & Neiva, 1912). Revista da Sociedade Brasileira de Medicina Tropical 45 (3): 313–317. https://doi.org/10.1590/S0037-86822012000300007

Rangel EF, Lainson R (2009) Proven and putative vectors of American cutaneous leishmaniasis in Brazil: aspects of their biology and vectorial competence. Memórias do Instituto Oswaldo Cruz 104 (7): 937–954. https://doi.org/10.1590/S0074-02762009000700001

Rangel EF, Lainson R, Carvalho BM, Costa SM, Shaw JJ (2018) Sand Fly vectors of American Cutaneous Leishmaniasis in Brazil. In: Rangel EF, Shaw JJ (Eds) Brazilian sand flies. Springer, Cham, Switzerland & Berlin, Germany, 341–380. https://doi.org/10.1007/978-3-319-75544-1_7

Shaw JJ, Rosa A, Ribeiro Cruz AC, Vasconcelos P (2018) Brazilian phlebotomines as hosts and vectors of viruses, bacteria, fungi, protozoa (excluding those belonging to the genus *Leishmania*) and nematodes. In: Rangel EF, Shaw JJ (Eds.) Brazilian sand flies. Springer, Cham, Switzerland & Berlin, Germany, 417–441. https://doi.org/10.1007/978-3-319-75544-1_9

Shaw JJ, De Faria DL, Basano SA, Corbett CE, Rodrigues CJ, Ishikawa EA, Camargo LM (2007) The aetiological agents of American cutaneous leishmaniasis in the municipality of Monte Negro, Rondônia state, western Amazonia, Brazil. Annals of Tropical Medicine and Parasitology 101 (8): 681–688. https://doi.org/10.1179/136485907x229103

Silva BQD, Afonso M, Freire LJM, Santana ALF, Pereira-Colavite A, Rangel EF (2022) Ecological aspects of the Phlebotominae fauna (Diptera: Psychodidae) among forest fragments and built areas in an endemic area of American visceral leishmaniasis in João Pessoa, Paraíba, Brazil. Insects 13 (12): 1156. https://doi.org/10.3390/insects13121156

SINAN (Sistema de Informação de Agravos de Notificação) (2023) Portal da Saúde. Doenças e Agravos de Notificação – 2007 em diante. DATASUS/Ministério da Saúde, Brazil. https://datasus.saude.gov.br/acesso-a-informacao/doencas-e-agravos-de-notificacao-de-2007-em-diante-sinan/. Acessed on: 2023-01-09.

Temponi AOD, Brito MG, Ferraz ML, Diniz SA, Silva MX, Cunha TN (2018) Ocorrência de casos de leishmaniose tegumentar americana: uma análise multivariada dos circuitos espaciais de produção, Minas Gerais, Brasil, 2007 a 2011. Cadernos de Saúde Pública 34 (2): e00165716. https://doi.org/10.1590/0102-311X00165716

Thomaz-Soccol V, Goncalves AL, Baggio RA, Bisetto A, Jr., Celestino A, Hospinal-Santiani M, de Souza A, Mychalizen M, Borges ME, Piechnik CA (2023) One piece of the puzzle: modeling vector presence and environment reveals seasonality, distribution, and prevalence of sandflies and *Leishmania* in an expansion area. One Health 17: 100581. https://doi.org/10.1016/j.onehlt.2023.100581

Uzcátegui YDVS, Vasconcelos Dos Santos T, Silveira FT, Ramos PKS, Dos Santos EJM, Póvoa MM (2020) Phlebotomines (Diptera: Psychodidae) from a urban park of Belém, Pará state, northern Brazil and potential implications in the transmission of American cutaneous leishmaniasis. Journal of Medical Entomology 57 (1): 281–288. https://doi.org/10.1093/jme/tjz153

Vilela ML, Zwetsch A, Silvas JDS (2018) Methods for capturing, processing and preserving Phlebotominae. In: Rangel EF, Shaw JJ (Eds.) Brazilian sand flies. Springer, Cham, Switzerland & Berlin, Germany, 443–466. https://doi.org/10.1007/978-3-319-75544-1_10

Vilela ML, Azevedo CG, Carvalho BM, Rangel EF (2011) Phlebotomine fauna (Diptera: Psychodidae) and putative vectors of leishmaniases in impacted area by hydroelectric plant, state of Tocantins, Brazil. PLoS ONE 6 (12): e27721. https://doi.org/10.1371/journal.pone.0027721